	ŧ
	Ħ
3	₹
_	3
0	
<u>_</u>	
_	Ξ
0	S

5

m

78

N.

07-10-626 Rec'n' PCT/PTO PTO-1390 Modified) U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFIC

TRANSMITTAL LETTER TO THE UNITED STATES

294-104 PCT/US

U.S. APPLICATION NO. (IF KNOWN, SEE 37 CFR

DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. 371 INTERNATIONAL APPLICATION NO.

INTERNATIONAL FILING DATE

TITLE OF INVENTION

PCT/NL00/00059

28 January 2000

PRIORITY DATE CLAIMED 29 January 1999

Composition Based on Cross-Linked Starch and Depolymerized Starch Suitable as Gelatine Replacement APPLICANT(S) FOR DO/EO/US Jakob Roelf WOLTJES, Heine Roelf MEIMA, and Pieter Lykle BUWALDA Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information: This is a FIRST submission of items concerning a filing under 35 U.S.C. 371. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371. 2. 3. This is an express request to begin national examination procedures (35 U.S.C. 371(f)). The submission must include itens (5), (6), (9) and (24) indicated below. The US has been elected by the expiration of 19 months from the priority date (Article 31). A copy of the International Application as filed (35 U.S.C. 371 (c) (2)) is attached hereto (required only if not communicated by the International Bureau). has been communicated by the International Bureau. is not required, as the application was filed in the United States Receiving Office (RO/US). ☐ An English language translation of the International Application as filed (35 U.S.C. 371(c)(2)). a.  $\square$ is attached hereto. b. 

has been previously submitted under 35 U.S.C. 154(d)(4). 7. Maximum Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371 (c)(3)) are attached hereto (required only if not communicated by the International Bureau). have been communicated by the International Bureau. have not been made; however, the time limit for making such amendments has NOT expired. have not been made and will not be made. An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)). ☐ An oath or declaration of the inventor(s) (35 U.S.C. 371 (c)(4)). 10. An English language translation of the annexes of the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371 (c)(5)). 11.  $\boxtimes$ A copy of the International Preliminary Examination Report (PCT/IPEA/409). 12. A copy of the International Search Report (PCT/ISA/210). Items 13 to 20 below concern document(s) or information included: 13 An Information Disclosure Statement under 37 CFR 1.97 and 1.98. 14. An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included. 15. □ A FIRST preliminary amendment. 16. A SECOND or SUBSEQUENT preliminary amendment. A substitute specification. 17. 18. A change of power of attorney and/or address letter. 19. A computer-readable form of the sequence listing in accordance with PCT Rule 13ter.2 and 35 U.S.C. 1.821 - 1.825. 20. A second copy of the published international application under 35 U.S.C. 154(d)(4). 21 A second copy of the English language translation of the international application under 35 U.S.C. 154(d)(4). 22. Certificate of Mailing by Express Mail 23. ☐ Other items or information:

**EXPRESS MAIL CERTIFICATE** 

07/09/01 Label No EL9220995 2805

Commissioner for Patents, by "EXPRESS MAIL Post O ASSANDRA LENS ( Page 1 of 2

PCTUS1/REV03

	in the
	1
	(I)
	00
	ű
	Sie
ş	grás
•	m
,	E.
	Sed.
	1
	N
	N
	O
	Ļį.

s. application in the state of	INTERNATIONAL APPLICATION PCT/NL00/00055				OCKET NUMBER PCT/US
4. The following fees are submitted:.				CALCULATIONS	PTO USE ONLY
ASIC NATIONAL FEE ( 37 CFR 1.492 (a) (1) -	(5)):				
☐ Neither international preliminary examination international search fee (37 CFR 1.445(a)(2) and International Search Report not prepared	) paid to USPTO I by the EPO or JPO	\$1	000.00		
	sared by the EPO of JPO	\$	860.00		
☐ International preliminary examination fee (3' but international search fee (37 CFR 1.445(a	i)(2)) paid to USP1U	:	5710.00		
International preliminary examination fee (3' but all claims did not satisfy provisions of Po	C1 Article 33(1)-(4)	!	690.00		
☐ International preliminary examination fee (3 and all claims satisfied provisions of PCT A	rticle 33(1)-(4)IATE BASIC FEE AMO		\$100.00 =	\$860.00	
			30	3000.00	
furcharge of \$130.00 for furnishing the oath or dec nonths from the earliest claimed priority date (37 G				\$0.00	
CLAIMS NUMBER FILED	NUMBER EXTRA		ATE		
otal claims 17 - 20 =	0		8.00	\$0.00 \$0.00	
ndependent claims 1 - 3 =	0	-	30.00	\$270.00	
Multiple Dependent Claims (check if applicable).	TANKE CALCULAT		<u> </u>	\$1,130.00	
TOTAL O	F ABOVE CALCULAT	IUNS		31,130.00	
Applicant claims small entity status. (See 37 C reduced by 1/2.	CFR 1.27). The fees indicated abo	ve are		\$0.00	
	SUB	TOTA	L =	\$1,130.00	
rocessing fee of \$130.00 for furnishing the Englishonths from the earliest claimed priority date (37)	th translation later than 2 CFR 1.492 (f)).	.0 [	□ 30 +	\$0.00	
	TOTAL NATIONA	L FEI	<u> </u>	\$1,130.00	
Fee for recording the enclosed assignment (37 CFF accompanied by an appropriate cover sheet (37 CFF)	R 1.21(h)). The assignment must R 3.28, 3.31) (check if applicab	be le).		\$0.00	
accompanies of an appropriate	TOTAL FEES ENCI	OSE	D =	\$1,130.00	
				Amount to be: refunded	\$
				charged	S
a. A check in the amount of \$1,	130.00 to cover the above fee	es is enc	losed.		
b. Please charge my Deposit Account A duplicate copy of this sheet is en	No in the an	nount of		to cover	the above fees.
c. The Commissioner is hereby author to Deposit Account No. 08-2	rized to charge any additional fee	s which s sheet is	may be re enclosed	equired, or credit any	overpayment
d. Fees are to be charged to a credit coinformation should not be included.	ard WARNING: Information on	this for	m may be	come public. Credit	card TO-2038.
NOTE: Where an appropriate time limit under 1.137(a) or (b)) must be filed and granted to res	. 37 CFP 1 494 or 1 495 has not	been m			
SEND ALL CORRESPONDENCE TO:		6		1/Ban	
		erc.	NATURE	y. v.	
Ronald J. Baron, Esq.					
Hoffmann & Baron, LLP					
Hoffmann & Baron, LLP 6900 Jericho Turnpike		Ro	ıald J. B	aron	
Hoffmann & Baron, LLP 6900 Jericho Turnpike Syosset, New York 11791		Roi NA	ME	aron	
Hoffmann & Baron, LLP 6900 Jericho Turnpike Syosset, New York 11791 Telephone: 516-822-3550		NA. 29,	ме 281		
Hoffmann & Baron, LLP 6900 Jericho Turnpike Syosset, New York 11791		NA. 29,	ме 281	ION NUMBER	
Hoffmann & Baron, LLP 6900 Jericho Turnpike Syosset, New York 11791 Telephone: 516-822-3550		Roi NA 29,	ме 281	ION NUMBER	

JOIS PROPRESSION OF JUL 2001

# Rec'd PCT/PTU 17 AUG ZUUI

## 09/889116 PATENT

## IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

plication No.: 09/889,116

Group No.: Unassigned

Inventors: Woltjes et al.

Examiner: Unassigned

Filed: July 9, 2001

Docket No.: 294-104 PCT/US

For: COMPOSITION BASED ON

Dated: August 15, 2001

CROSS-LINKED STARCH AND DEPOLYMERIZED STARCH

SUITABLE AS GELATINE REPLACEMENT

Assistant Commissioner for Patents

Washington, DC 20231

I hereby certify this correspondence is being deposited with the U.S. Postal Service as first class mail, postpaid in an envelope, addressed to: Asst Commissioner for Patents, Washington, D.C.

20231 on August 15, 2001 Signature

PRELIMINARY AMENDMENT

Sir:

In order to place the application in proper form for examination, applicants

hereby amend the application as follows.

#### IN THE SPECIFICATION:

Please amend the specification as follows:

Page 1, after the title of the invention, insert the following:

-- BACKGROUND OF THE INVENTION --

Page 3, after line 32, insert the following:

--SUMMARY OF THE INVENTION--

Page 12, line 1, change "Detailed description" to the following:

-- DESCRIPTION OF THE PREFERRED EMBODIMENTS--

#### IN THE CLAIMS:

### Please delete claims 9 to 15.

(NOTE: For the examiner's convenience, applicants are listing all of the claims and indicating the amended claims.)

## Please amend the claims as follows:

- (Amended) A starch composition comprising a first fraction comprising cross-linked starch and at least a second fraction comprising depolymerized starch.
- A composition according to claim 1 wherein said first fraction comprises starch cross-linked by using sodium trimetaphospate, phosphorus oxytrichloride or adipic anhydride.
- (Amended) A composition according to claim 1 wherein said first fraction further comprises stabilized starch.
- (Amended) A composition according to claim 3 wherein said first fraction is stabilized by hydroxyalkylation.
- (Amended) A composition according to claim 1 wherein said second fraction comprises depolymerized starch obtained by oxidation.
- (Amended) A composition according to claim 1 wherein said second fraction further comprises stabilized starch.
- (Amended) A composition according to claim 6 wherein said second fraction is stabilized by acetylation.
- (Amended) A composition according to claim 1 wherein said starch is derived from potato.

- 9. (Deleted)
- 10. (Deleted)
- 11. (Deleted)
- 12. (Deleted)
- 13. (Deleted)
- 14. (Deleted)
- 15. (Deleted)

#### Please add the following claims:

- 16. A method for preparing a thickened foodstuff comprising mixing a starch composition comprising a first fraction comprising cross-linked starch and at least a second fraction comprising depolymerized starch with a water-based liquid.
- A method according to claim 16 wherein said foodstuff comprises confectionery.
- 18. Foodstuff comprising a starch composition comprising a first fraction comprising cross-linked starch and at least a second fraction comprising depolymerized starch.
- A foodstuff according to claim 18 wherein the use of said composition at least partly replaces the use of gelatine in said foodstuff.
- Confectionery comprising a starch composition comprising a first fraction comprising cross-linked starch and at least a second fraction comprising depolymerized starch.

## MARKED-UP VERSION SHOWING ALL CHANGES MADE

### IN THE CLAIMS:

## Please amend the claims as follows:

- (Amended) A starch composition comprising a first fraction comprising cross-linked starch and at least a second fraction comprising depolymerised depolymerized starch.
- A composition according to claim 1 wherein said first fraction comprises starch cross-linked by using sodium trimetaphospate, phosphorus oxytrichloride or adipic anhydride.
- 3. (Amended) A composition according to claim 1 or 2 wherein said first fraction further comprises stabilised stabilized starch.
- 4. (Amended) A composition according to claim 3 wherein said first fraction is stabilised stabilized by hydroxyalkylation.
- (Amended) A composition according to anyone of elaims 1 to 4 claim 1 wherein said second fraction comprises depolymerised depolymerized starch obtained by oxidation.
- (Amended) A composition according to any one of claims 1 to 5 claim 1 wherein said second fraction further comprises stabilised stabilized starch.
- (Amended) A composition according to claim 6 wherein said second fraction is stabilised stabilized by acetylation.
- (Amended) A composition according to anyone of claims 1-7 claim 1 wherein said starch is derived from potato.

- (Deleted) Use of a starch composition according to anyone of claims 1 to 8 in the preparation of a foodstuff.
- 10. (Deleted) Use according to claim 9 wherein the use of said composition at least partly replaces the use of gelatine in said foodstuff.
- 11. (Deleted) Use according to claim 9 or 10 wherein said foodstuff comprises confectionery.
- 12. (Deleted) A method for preparing a thickened foodstuff comprising mixing a starch composition according to anyone of claims 1 to 8 with a water-based liquid.
- 13. (Deleted) A method according to claim 12 wherein said foodstuff comprises confectionery.
- 14. (Deleted) Foodstuff comprising a starch composition according to anyone of claims 1 to 8:
- (Deleted) Confectionery comprising a starch composition according to anyone of claims 1 to 8.

## Please add the following claims:

- 16. A method for preparing a thickened foodstuff comprising mixing a starch composition comprising a first fraction comprising cross-linked starch and at least a second fraction comprising depolymerized starch with a water-based liquid.
- 17. A method according to claim 16 wherein said foodstuff comprises confectionery.

18. Foodstuff comprising a starch composition comprising a first fraction comprising cross-linked starch and at least a second fraction comprising depolymerized starch.

- 19. A foodstuff according to claim 18 wherein the use of said composition at least partly replaces the use of gelatine in said foodstuff.
- 20. Confectionery comprising a starch composition comprising a first fraction comprising cross-linked starch and at least a second fraction comprising depolymerized starch.

#### CONCLUSION

Applicants respectfully submit that the application is now in proper form for examination and request early examination.

Dated: August 15, 2001

Kevin E. McDermott Registration No.: 35,946 Attorney for Applicants

HOFFMANN & BARON, LLP 6900 Jericho Turnpike Syosset, New York 11791 (516) 822-3550

139160\_1.DOC

20

25

30

09/889116

# COMPOSITION BASED ON CROSS-LINKED STARCH AND DEPOLYMERIZED STARCH SUITABLE AS GELATINE REPLACEMENT

The invention relates to hydrocolloids that are used as thickening, binding, gelling or stabilising agents in the food industry.

In the food industry hydrocolloids are in general used to thicken (gel, bind, stabilise) water-based foodstuffs. Gelatine is a popular hydrocolloid, which, contrary to other hydrocolloids which are mainly of a polysaccharide nature, is a protein. Gelatine is derived from animal slaughter offal, such as skins and bones, by hydrolysis of insoluble collagen into soluble gelatine. Collagen is the major structural component of white tissue fibres and present in all tissues and organs of animals where it constitutes almost 30% of total protein content.

Gelatine is used in a great number of food applications, where it is desired because it has a number of characteristics that are superior over other hydrocolloids used in the food industry. It is used for example as a thickening or gelling agent in jellied products such as confectionery and aspic type of foods; as a stabiliser and thickener in ice cream and icings, as emulsifier and thickener in dressings, desserts and sauces, as thickener in syrups and soups, as binder or thickener in general and as fining agent. For example, gelatine is used in gum and jelly products, such as wine gums, as gelling agent to give the end product an elastic, gummy structure. In particular, gelatine is superior over other thickening (gelling and binding) agents for the clarity and elasticity it renders to the food product.

The use of gelatine (or hydrolysed collagen) in the food industry, however, has recently been criticised because of its animal origin. Traditionally, gelatine containing foodstuff has been avoided by vegetarian

15

25

30

35

consumers and by consumers whose religion teaches to avoid animal derived products like gelatine. These traditional gelatine avoiding consumers were in general satisfied with buying products that contained less superior binding agents to accommodate their vegetarian and/or religious preferences. More recently, however, the general consumer, albeit not bound by vegetarian or religious preferences, is also shifting to a preference for foodstuff wherein gelatine is replaced by another agent. Said shift in the preference of the general public is mainly understood to have been initiated by the recent occurrence of prion diseases such as seen with mad cow disease, and by concern that these prion diseases may infect humans when proteinaceous food of animal origin is eaten.

The prion diseases bovine spongiform encephalopathy (BSE) and scrapie of cattle and sheep, respectively, are fatal neurodegenerative diseases caused by prion proteins and are characterised by a long incubation period. In humans Creutzfeldt-Jakob disease (CJD), Gerstmann-Sträussler-Scheinker syndrome (GSS) and fatal familial insomnia belong to this category of transmissible spongiform encephalopathies (TSEs). Although scrapie, the prototype of the family of TSEs, in sheep and goats has been known for over 200 years and has been diagnosed world-wide, it is only since 1986 that BSE has been described in cattle in the UK. By January 1998, there had been 170,259 confirmed cases of BSE in Great Britain and there may exist a great number of cases of not yet overt cases of BSE. BSE apparently emerged because scrapie contaminated sheep offal, via meat and bone meal had been included in cattle feeding-stuff, and newly infected cattle material was then recycled and eaten by susceptible cattle. Brain homogenates from cows with BSE produce a characteristic pattern of brain lesions in mice. This is identical to the pattern elicited by brain

20

35

tissue from individuals who recently have died from newvariant Creutzfeldt-Jakob disease. Up to now, this variant has caused the death of 35 young Britons and one Frenchman.

There is also concern that the BSE strain that seems to be transmissible to humans may have infected sheep, where it could produce a disease hardly distinguishable from scrapie. Sheep BSE may be a threat to human health, although scrape by itself seems not to transmit to humans. Indeed, BSE agent has been transmitted experimentally to sheep by the oral route and thus could have the potential to infect sheep under field conditions.

Thus far, the only known cause of prion disease is an abnormal form of the normal prion protein called aberrant prion protein. Said aberrant prion protein is mainly characterised by its resistance to proteolytic hydrolysis, it is typically quite resistant against treatment with high or low pH, and generally only looses its infectivity after prolonged treatment under high temperature.

Although most governments in Western society have taken strict measurements to alleviate public concerns related to mad cow disease, for example by strictly

25 banning the use of animal products derived from animals with prion disease in the food industry, public concerns related to using the protein derived gelatine still exist, and seem to be growing. Consequently, among the general public lives a growing desire to consume nongelatine derived foodstuff, that, however, has similar or comparable superior characteristics as the traditionally gelatine comprising foodstuffs have.

It is an object of the present invention to provide a non-protein hydrocolloid that has suitable characteristics to serve as gelatine replacement in

WO 00/44241 PCT/NL00/00059

foodstuff of varied nature or that can be used to prepare new types of foodstuff.

The invention provides a starch composition, comprising a first fraction comprising cross-linked starch and at least a second fraction comprising depolymerised starch. Said composition is suitable for example to serve as gelatine replacement in foodstuff of varied nature, and is preferably suitable for use in 10 foodstuff such as confectionery. Starches suitable for use in a composition according to the invention are for example chosen from maize, wheat, barley, rice, triticale, rice, millet, tapioca, arrow root, banana, potato, sweet potato starches or from high amylose starches like amylomaize, wrinkled pea starch, mung bean starch or from amylopectin rich starches like waxy maize, waxy barley, waxy wheat, waxy rice, amylopectin potato, amylopectin tapioca, amylopectin sweet potato or amylopectin banana starch. Amylopectin starches may be 20 derived from plants that selectively produce amylopectin such as waxy cereals or amylose-free potato mutants and/or genetically modified plant varieties such as potatoes modified to selectively produce amylopectine. Cross-linked starch in general is a modified starch in which cross-links between starch macromolecules have been formed by means of bifunctional or polyfunctional chemical reagents, and results in the formation of large complexes of starch molecules with high molecular weight. Cross-links can for example be formed between amylose molecules or between amylopectine molecules, or between amylose and amylopectine molecules in the starch. Although not preferred from the viewpoint of ease of production, a first fraction according to the invention can of course also be partly depolymerised, or be modified in any other way, and a second fraction according to the invention can also be partly crosslinked, or be modified in any other way, if so desired.

20

25

35

Cross-linking starch in itself is a method known in the art and various agents are known. Examples are: epichlorohydrin, sodium trimetaphosphate, phosphorous oxychloride (POCl3), adipic anhydride, or other reagents with two or more halogen, halohydrin or epoxide groups or combinations which all can be used as cross-linking agents. Preferred are distarch phosphates and distarch adipates. A cross-linked or cross-bonded starch may for example be cross-bonded by 0.003 to 0.024% of adipic anhydride, preferably by 0.01 to 0.03%. Prior to crossbonding by adipic anhydride the starch may be treated with hydrogen peroxide and/or peracetic acid. Preferably with a quantity which corresponds to 0.001% to 0.045% of active oxygen, most preferably to 0.005 to 0.045%. A 15 distarch phosphate may for example be cross-bonded by sodium trimetaphosphate up to such a degree that the residual phosphate is no more than 0.14% for a potato starch or 0.04% for other starches. Preferably the starch is cross-bonded with 0.01% to 0.25%, most preferably with 0.025 to 0.15% of sodium trimetaphosphate, under conditions known to the artisan. Of course it is always possible for the artisan to find conditions in which the reactants react with a very low yield, outside of the preferred conditions resulting in a starch with desired properties. A distarch phosphate may as well be crossbonded with phosphorous oxychloride up to such a degree that the residual phosphate is not more than 0.14% for a potato starch or 0.4% for other starches. Preferably the starch is cross-bonded with 0.00010 % to 0.01% of phosphorous oxychloride, under conditions known to the artisan. Of course it is always possible for the artisan to find conditions in which the reactants react with a very low yield, outside of the preferred conditions resulting in a starch with the desired properties.

Depolymerised starch is in general obtained by a reduction in the degree of polymerisation of the starch chains by physical, chemical or enzymatic action, and results in general in a reduction of the size (molecular

weight) of the starch molecules. Depolymerisation by acid hydrolysis of starch can be accomplished in many ways known to the artesian. Preferred enzymatic hydrolysis, acid catalysed hydrolysis using organic or mineral acid and dry roasting of starches under acidic conditions (dextrination). In an embodiment of the invention the starch is hydrolysis in a slurry process using hydrochloric or sulphuric acid. During hydrolysis with acid the molecular weight of the starch by hydrolytic cleavage of the acetal function of the starch. 10 Depolymerisation by oxidation of starch can also be accomplished an large number of ways. Known oxidation agents are sodium hypochlorite, calcium hypochlorite, sodium and potassium permanganate and hydrogen peroxide. 15 Preferred is the hypochlorite oxidation of starch which is still one of the most useful reactions for the derivatisation of starches. Oxidised starches are applicable in a wide range of applications such as paper manufacture, adhesives, textile industry and food. 20 Hypochlorite is a relatively cheap oxidation agent. During oxidation with hypochlorite different oxidation reaction may take place, of which the hydrolysis of the glucosidic bonds is the most important. Together with a decrease in molecular weight of the starch molecule also functional groups such as carbonyl and carboxyl groups are introduced. The instability of the glucosidic bonds is influenced by the presence of carbonyl or carboxyl groups. The course of an oxidation reaction is primarily controlled by the amount of hypochlorite, the pH, the temperature and the presence of a catalyst. An overview

30 of the most important reaction parameters is depicted in article by J. Potze and P. Hiemstra in Starch; volume 15, page 217-225 (1963). The decrease in molecular weight, and the amount of functional groups is direct 35 proportional to the amount of hypochloride. The oxidation

rate is very sensitive for the pH during oxidation. The

highest reaction rates are obtained at neutral pH (pH about 7), the reaction rate decreases with increasing pH. The carbonyl content of oxidised starches is also controlled by the pH. At neutral pH the highest carbonyl contents are obtained. Due to this high amount of carbonyl groups, also a high amount of hydrolyses of the glucosidic bonds is observed at neutral pH, resulting in a decrease in viscosity of the resulting oxidised starch derivative. Increasing the oxidation pH results in a steady decrease in both carbonyl content and hydrolysis of the glucosidic bonds. Unfortunately, a high amount of carbonyl groups leads to viscosity instability of the oxidised starch. The introduction of carboxyl groups give rise to an increase in viscosity stability of the 15 oxidised starch. Here too the amount of carboxyl groups is controlled by the pH, with a maximum at pH 9. Hypochlorite oxidations are mainly performed in slurry processes. The degradation of starch can be monitored by measuring the intrinsic viscosity. The intrinsic 20 viscosity of depolymerised granular products lies between 0.2 and 2.5 dL/g, preferably between 0.3 and 1.5, mostly preferred between 0,4 and 1.2.

Surprisingly, by combining said fractions, each having distinct characteristics, a starch composition is 25 obtained that demonstrates characteristics comparable to characteristics of gelatine, or even improving those of gelatine, most notably those relating to clarity and elasticity of the foodstuff when said composition is used as binding, thickening or gelling agent. Combining said two fractions according to the invention imparts said desired characteristics to the starch composition, making it suitable to use the composition as provided by the invention in the food industry to partly or even fully replace gelatine and to prepare new types of foodstuff. Said composition comprising said two fractions can for 35 example be obtained by mixing separate starch fractions,

a cross-linked fraction and a depolymerised fraction, in the desired ratio to obtain a starch composition as provided by the invention, however, it is also possible to obtain said composition by partial cross-linking and partial depolymerisation of a starch. However, a mixture of at least two fractions is preferred in the light of obtaining a starch composition according to the invention that is adjusted to for example the level of elasticity and/or clarity required of the foodstuff.

By virtue of the diverse fractions, said starch composition demonstrates characteristics that are close to or even improved over those of gelatine. The ratio at which the fractions shall be used are easily determined by mixing and testing the thus obtained compositions for desired properties. Suitable ratios of first fraction to second fraction (weight%:weight%) vary for example from 5:95 to 95:5, preferably from 10:90 to 90:10, more preferably from 25:75 to 75:25. For example, soft food products, such as soft sugar confectionery in which gelatine is traditionally used most, desire most elasticity, and by adjusting the ratio of the two fractions in the starch composition as provided by the invention, the desired clarity and/or elasticity can be obtained. Also, in hard confectionery products, where traditionally hydrolysed collagen is used in mixtures to replace gums, such as gum arabic, clarity and/or elasticity are now obtained using a starch composition according to the invention. In general when more elasticity is required, one increases the cross-linked fraction, and when more gelling and/or clarity is desired one increases the depolymerised fraction.

In a preferred embodiment, the invention provides a starch composition comprising a first fraction comprising cross-linked starch and at least a second fraction comprising depolymerised starch wherein said first fraction is cross-linked by using sodium trimetaphospate,

15

phosphorus oxytrichloride or adipic anhydride, using for example a method known in the art. These cross-linking agents are most suitable for use in the food industry.

Preferred are cross-linked starch acetates having an acetyl content which corresponds to a DS or degree of substitution of 0.001 to 0.2, preferably from 0.03 to 0.092, most preferably from 0.05 to 0.092. The term DS used herein indicates the average number of sites per anhydroglucose unit of the starch molecule in which there are substituent groups.

Even more preferred are cross-linked hydroxypropylated starches having a hydroxypropyl content which corresponds to a DS of 0.001 to 0.3, preferably, 0.03 to 0.21, most preferably 0.06 to 0.21.

In a further preferred embodiment the invention provides a starch composition according to the invention wherein said first fraction is also stabilised. Stabilisation in general is done by methods known in the art, such as by treatment with acetic anhydride or vinyl acetate, or comparable agents, but for the purpose of gelatine 20 replacement a preferred embodiment is a composition according to the invention wherein said first fraction is stabilised by hydroxyalkylation, for example by hydroxypropylation. Stabilisation by hydroxyalkylation of starch is for example obtained with reagents containing a

25 halohydrin, or an epoxide group as reactive site. The addition of hydroxypropyl groups is generally performed in aqueous suspensions of starch using propylene oxide, under alkaline conditions. Cross-bonding and/or stabilising reagents are reacted with starch under 30 alkaline conditions. Suitable alkali materials are: sodium hydroxide, potassium hydroxide, ammonium hydroxide, magnesium hydroxide, sodium carbonate and

hydroxides and carbonates, most preferred are sodium hydroxide and sodium carbonate. Sometimes salts are added as to prevent swelling under alkaline reaction conditions. Preferred are sodium chloride and sodium

trisodium phosphate. Preferred are the alkali metal

sulphate.

In a further embodiment, the invention provides a starch composition comprising a first fraction comprising cross-linked starch and at least a second fraction comprising depolymerised starch wherein said depolymerisation is for example achieved by acid or enzymatic hydrolysis. Preferred for the purpose of gelatine replacement, however, is a composition according to the invention wherein said second fraction is obtained by oxidation of starch, such as by treatment with sodium hypochlorite or hydrogen peroxide.

In a further preferred embodiment the invention provides a starch composition according to the invention wherein said second fraction is also stabilised. 15 Stabilisation in general is done by methods known in the art, such as by hydroxyalkylation or by acetylation with acetic anhydride, vinyl acetate or comparable agents. For the purpose of gelatine replacement a preferred embodiment is a composition according to the invention wherein said second fraction is stabilised by 20 acetylation. Stabilisation by acetylation is performed using acetic anhydride or vinyl acetate. Other stabilisation reagents are for example succinic anhydride, 1-octenyl succinic anhydride, sodium 25 tripolyphosphate, potassium orthophosphate, sodium orthophosphate or orthophosphoric acid. In yet another embodiment, said second fraction is also cross-linked, for example by a suitable method as listed above, to provide even more elasticity (chewiness).

30 The invention also provides use of a starch composition according to the invention in the preparation of a foodstuff, preferably partly or fully replacing gelatine in said foodstuff. By using a composition according to the invention, it is possible to replace 35 gelatine for mote than 50%, up to 80% or even 100%, depending on the requirements of the customer. It is thus now possible to reduce gelatine content and select and

20

WO 00/44241 PCT/NL00/00059

use minimal quantities of those gelatine batches that are absolutely prion protein free, or to fully replace animal derived thickeners, such as hydrolysed collagen, or gelatine, that may be derived from slaughter offal comprising aberrant prion protein. In a preferred embodiment, said foodstuff comprises confectionery, for example sugar confectionery such as hard or soft sugar confectionery, lozenges or dragees, or confectionery for diabetics wherein the sugar is replaced by artificial sweeteners.

The invention furthermore provides a method for preparing a thickened (thickening herein also called gelling, stabilising or binding) foodstuff comprising mixing a starch composition according to the invention with a water-based liquid. Such a water-based liquid can for example be water, milk or another dairy product, a stock or bouillon, a sugar solution, a beverage or another water-based liquid food component known in the art. Of course, said method allows for the additional use of other ingredients, of which many are known in the art. Traditionally, gelatine was a first choice in thickening such liquids, however, due to consumer preference, a method to prepare non-delatine foodstuffs is desired.

Furthermore, the invention provides foodstuff

comprising a starch composition comprising a first
fraction comprising cross-linked starch and at least a
second fraction comprising depolymerised starch. Such
foodstuff can for example be characterised by a clarity
and/or elasticity that in general lives up to consumers

expectations relating to gelatine comprising foodstuffs.
In a preferred embodiment the invention comprises
confectionery, as further exemplified in the detailed
description herein without limiting the invention.

15

Detailed description

General overview of confectionery production

Sugar-based products, or sugar confectionery, can be divided into the following groups.

- Hard sugar confectionery (fruit drops, clear mints,
   barley sugars, and bonbons)
  - Soft sugar confectionery (gums, pastilles, jellies, chewing gums, gelées, liquorice)
  - 3. Lozenges (sheeted/pressed confectionery)
  - 4. Dragees (coated confectionery).

In sugar confectionery starch products, natural gums and proteins are traditionally used as binding and/or gelling agents. There are two main types of soft sugar: gummy products like fruit gums, liquorice gums, pastilles, etc.

and elastic products like wine gums, jellies, etc. A binding and/or gelling agent itself has specific properties, which influences the properties of the final product. For soft sugar confectionery applications gelatine is traditionally used as gelling agent, as it

imparts a desired elasticity to the end product. In soft sugar confectionery thin boiling starches are commonly used to replace gelatine in part, however replacement of gelatine with thin boiling starches leaves the customer with products having a low acceptance as they have little

30 elasticity.

The application of gelatine and mixtures of gelatine and thin boiling starches in soft confectionery is a well-known process. For instance in "Sugar Confectionery Manufacture" edited by E.B. Jackson, Blackie and Son, London 1990 an overview of different process equipment

and product recipes are given. Gums, pastilles and

15

20

25

WO 00/44241 PCT/NL00/00059

jellies can be processed using a variety of techniques, batch cooking, direct cooking (jet cooker), indirect cooking or extrusion cooking. Batch process proceeds through atmospheric cooking in open pans. Colouring, flavouring and moulding and drying in starch moulds follows cooking. Direct cooking comprises two parts: cooking and a vacuum cooling part. The confectionery mass is pumped into the continuous (jet)cooker where it is heated with steam. The pressure of the steam determines the cooking temperature. The slurry is then pumped into the vacuum cooler, where it is cooled and all air removed. When the required concentration is reached, colours and flavouring can be added. The slurry can then be formed.

The present invention among others relates to the partial or full replacement of gelatine in confectionery. This can be for example be achieved by a using a starch composition comprising a combination of a thin boiling starch and cross-linked and stabilised starch. In one embodiment of the present invention the thin boiling starch is prepared by oxidation or acid degradation of starch. The starch may be oxidised by sodium hypochlorite or by hydrogen peroxide. The acid degradation may be performed with hydrochloric acid or sulphuric acid in slurry process.

Alternatively, the acid degradation can be achieved through a dextrination process using hydrogen chloride or sulphuric acid applying heat. The cross-bonding can be performed with sodium trimetaphosphate, phosphorus oxytrichloride or adipic anhydride using procedures known to the artesian. The stabilisation can be performed with propylene oxide, acetic anhydride or vinyl acetate. A thin boiling starch is for example stabilised and oxidised or stabilised acid degraded.

Analytical procedures

Sensory evaluation

- 5. The confectionery was tested by a trained panel for a number of sensor features. The products were evaluated on a sensoric scale from 1 to 5, and a general acceptance scale from 1 to 10.
- 10 Testing the end products for the presence of cross-bonded starch derivatives

In order to determine the cross-linked starch in confectionery a sedimentation test was carried out.

15 Therefor 20 gram wine gum was dissolved in a potassium dihydrogen phosphate/ sodium hydroxide buffer at pH at 90 °C to give a solution of approx. 20%. The solution was cooled to room temperature and the amount of dry substance was measured with a refractometer. Then this solution was

20 diluted to a 5 % solution with deminarelised water. 100 ml solution was allowed to stand for 24 hours in a measuring cylinder. Afterwards the amount of sedimentation was estimated. The cross-linked starch could be identified under a microscope also.

25

Intrinsic viscosity (IV)

The intrinsic viscosity is determined in a known manner with a Ubbelohde viscosity meter with 1 M sodium hydroxide as solvent and expressed in g/dl.

Example 1

The preparation of cross-linked hydroxypropylated starch

35

A 39 % starch slurry is prepared. To this slurry sodium

sulphate (100 g/kg) and sodium hydroxide (7.5 g/kg starph) as 4.4 % solution are added. The temperature was raised to 35 °C and POCl3 is added (15 to 200  $\mu L/kg$  depending on the degree of cross-linking). Next

5 propylene oxide (DSmax = 0.225) is introduced and the reaction was allowed to proceed 20-24 hours. The slurry was neutralised with sulphuric acid to pH 5-6 and washed and dried using conventional means known to the art. The starch used are potato starch and tapioca starch.

10

Example 2

## The preparation of cross-linked acetylated starch

15 A 39 % starch slurry is prepared of 1 kg of potato starch. To this slurry 25 g of sodium chloride and sodium hydroxide (7.5 g/kg starch) as 4.4 % solution are added. The temperature was raised to 35 °C and sodium trimetaphosphate (400 or 600 mg depending on the degree of cross-linking) is added. The reaction is allowed to proceed for 6 hours. The slurry was neutralised with sulphuric acid to pH 8.5 and 61 g of acetic acid is added drop wise. After the decline of reaction the suspension is neutralised with sulphuric acid to pH 5-6 and washed and dried using conventional means known to the art.

Example 3

30

## The preparation of adipylated, acetylated starch

The adipate/acetate of amylopectin potato starch was prepared using a 1:20 mixture. 810 gram dry starch was suspended in water to 39 % suspension. The pH was adjusted to 8.5 with a 4.4 % (w/w) sodium hydroxide solution and 1.3 ml of 30 % of H2O2 solution were added. Then 6 g of 1:20 adipic anhydride reagent was added drop

wise keeping the pH at 8.5 with a 4.4 %(w/w) sodium hydroxide solution. Afterwards 41 grams of acetic anhydride was added drop wise to a pH 8.5. The acetic anhydride reaction was performed in approx. 1.5 hours.

5 After the acetic anhydride dosage the suspension was stirred for 10 minutes at pH 8.5. Then the suspension was neutralised with sulphuric acid to pH 5.5 and the final product was dewatered, washed and dried using methods known to the art.

10

Example 4

## The preparation of acid-degraded starch

5 A 39 % starch slurry is prepared of 1 kg of potato starch and water. To this slurry 80 ml of 10 N sulphuric acid are added. The temperature was raised to 45 °C. The reaction is allowed to proceed 17 hours. The slurry is neutralised with sodium hydroxide to pH 5-6 and washed and dried using conventional means known to the art.

Example 5

## The preparation of oxidised starch

25

Starch was oxidised in aqueous suspension with sodium hypochlorite using sodium hydroxide as a catalyst. Thus 810 gram dry starch was suspended in water to 39 % suspension. The temperature of the suspension was 30 adjusted at 35 °C. Sodium hypochlorite was added, amounts corresponding with 10 grams of active chlorine and at the same time sodium hydroxide was added, as a 4.4 % (w/w) solution, adjusting the pH to a value of 11. During the reaction the pH is kept at 11. When there was no active chlorine detectable the suspension was bleached with 5 ml sodium hypochlorite and after 1 minute the

excess of chlorine was removed using sodium hydrogen sulphite.

Afterwards the suspension was neutralised with sulphuric acid to pH 5.5 and the final product was dewatered, washed and dried using methods known to the art.

Example 6

## The preparation of oxidised acetylated starch

10

15

20

25

Potato starch was oxidised in aqueous suspension with sodium hypochlorite using sodium hydroxide as a catalyst. Thus 810 gram dry starch was suspended in water to 39 % suspension. The temperature of the suspension was adjusted at 35 °C. Sodium hypochlorite was added, amounts corresponding with 10 grams of active chlorine and at the same time sodium hydroxide was added, as a 4.4 %(w/w) solution, adjusting the pH to a value of 11. During the reaction the pH is kept at 11. When there was no active chlorine detectable the suspension was bleached with 5 ml sodium hypochlorite and after 1 minute the excess of chlorine was removed using sodium hydrogen sulphite. Afterwards the suspension was neutralised with sulphuric acid to pH 8.5. The temperature was adjusted to 25 °C and the suspension was acetylated using 36 g of acetic anhydride at a constant pH value of 8.5. The acetic anhydride reaction was performed in approx. 1.5 hours. After the acetic anhydride dosage the suspension was stirred for 10 minutes at pH 8.5. Then the suspension was neutralised with sulphuric acid to pH 5.5 and the final product was dewatered, washed and dried using methods known to the art. According to the examples 1-6 the following starches were prepared.

Table 1.

product	starch	modification	amount of
			cross-linker
1	potato	oxidation	
2	tapioca	hydroxypropylation/	85 µL/kg
		cross-linking	
3	potato	hydroxypropylation/	15 μL/kg
		cross-linking	
4	potato	hydroxypropylation/	40 μL/kg
		cross-linking	
5	potato	acetylation/	100 μL/kg
		cross-linking	
6	potato	acetylation/	400 mg/kg
		cross-linking	
7	potato	acetylation/	600 mg/kg
		cross-linking	]
8	amylopectin	adipylation/	6 g/kg
	potato	acetylation	
9	potato	acid	
		degradation	ļ
10	potato	oxidation/	
		acetylation	

#### Example 7

### Preparation of wine gum using batch process

A starch suspension is prepared using a starch: water ratio of 1: 1.5 Into a batch-cooker (open pan) hot water is filled. Glucose-syrup (DE 42; Dormamix 42/82, Pfeiffer & Langen) is added and stirring is started. The

- Pfeiffer & Langen) is added and stirring is started. The cooker is heated and sugar is added while stirring. The heating continued until the mixture starts to boil. At this moment the starch suspension is added slowly, while stirring. Boiling is maintained until the desired dry solid is reached (74 -76 %)
- The mixture is cooled to 80 °C and colour, flavour and citric acid are added. The cooked solution is moulded into shapes in moulding powder. The resulting are dried for 24 hours at 50 °C.
- 20 Example 8

#### Preparation of wine gum using direct cooking

A premix is made of a recipe containing sugar/glucose

syrup (DE42; Dormamix 42/82, Pfeiffer & Langen), starches
and water. This premix is cooked at 1300C in a

continuous cooker system from Vomatec. The cooked
solution is cooled by applying vacuum. Colour, flavour
and citric acid are added to the cooled solution. This

solution is moulded into shapes in moulding powder. The
moulded products are dried and subsequently evaluated.

According to example 8 the following mixtures were prepared and moulded into shapes.

35

Table 2.

WO 00/44241

Exp. nr	A	В	С	D	E	F	G	Н	I	J	к	L
Starch comp.												
1	9	9	9	9	9	9	9		8		12	
2	3									$\top$		
3		3				$\top$	T	3		1		3
4			3			T						
5				3								1
6					3	T				1		1
7						3				$\top$		
8							3					1
9								9		8		
gelatine*									4		T	
gelatine*										4		
10												9
sugar	34	34	34	34	34	34	34	34	30	30	34	34
Slucose-syrup	34	34	34	34	34	34	34	34	38	38	34	34
ater	20	20	20	20	20	20	20	20	20	20	20	20

gelatine 240 bloom

The resulting products were evaluated. The results are summarised in table 3.

Table 3.

Exp.	elast.	clarity	general
A	4	3	7
В	4	3	6
С	4	3	7
D	4	2	3
E	3	3	4
F	3	2	3
G	3	3	3
Н	3	3	3
I	4	3	7
J	4	3	7
K	1	1	2
L	5	4	8

## Example 9

5

## Preparation of liquorice

	Product 1	9%
	Product 3	3%
10	Sugar	28%
	Glucose-syrup DE 42	32%
	Ammonium chloride	4%
	Liquorice powder	4%
	Water	20%

15 The resulting liquorice had a gummy, elastic texture

### CLAIMS

- 1. A starch composition comprising a first fraction comprising cross-linked starch and at least a second fraction comprising depolymerised starch.
- A composition according to claim 1 wherein said first fraction comprises starch cross-linked by using
- sodium trimetaphospate, phosphorus oxytrichloride or adipic anhydride.
- A composition according to claim 1 or 2 wherein said first fraction further comprises stabilised starch.
- 4. A composition according to claim 3 wherein said first fraction is stabilised by hydroxyalkylation.
  - A composition according to anyone of claims 1 to 4 wherein said second fraction comprises depolymerised starch obtained by oxidation.
- 15 6. A composition according to any one of claims 1 to 5 wherein said second fraction further comprises stabilised starch.
  - 7. A composition according to claim 6 wherein said second fraction is stabilised by acetylation.
- 20 8. A composition according to anyone of claims 1-7 wherein said starch is derived from potato.
  - 9. Use of a starch composition according to anyone of claims 1 to 8 in the preparation of a foodstuff.
    - 10. Use according to claim 9 wherein the use of said
- 25 composition at least partly replaces the use of gelatine in said foodstuff.
  - 11. Use according to claim 9 or 10 wherein said foodstuff comprises confectionery.
- 12. A method for preparing a thickened foodstuff 0 comprising mixing a starch composition according to anyone of claims 1 to 8 with a water-based liquid.
  - 13. A method according to claim 12 wherein said foodstuff comprises confectionery.

14. Foodstuff comprising a starch composition according to anyone of claims 1 to 8.

15. Confectionery comprising a starch composition according to anyone of claims 1 to  $8. \,$ 

1-00

## **Power of Attorney**

As a named inventor, I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and transact all business in the Patent and Trademark Office connected therewith.

Attorney Registration Numb	er
Charles R. Hoffmann         24.102           Ronald J. Baron         29,281           Gerald T. Bodner         30,449           A. Thomas Kammer         28,226           Irving N. Feit         28,601           Alan M. Sack         31,874           Algis Anilionis         36,995           Gregory W. Bachmann         41,593           Anthony E. Bennett         40,910           James F. Harrington         P-44,741           Glenn T. Henneberger         36,074           Richard LaCava         41,135           Kevin E. McDermott         35,946           Robert C. Morriss         42,910           Samir R. Patel         P-44,998           R. Glenn Schroeder         34,720           Susan A. Sipos         43,128           Roderick S.W. Turner         38,639           Steven T. Zuschlag         43,309	(19)

I hereby authorize them or others whom they may appoint to act and rely on instructions from and communicate directly with the person/organization who/which first sends this case to them and by whom/which I hereby declare that I have consented after full disclosure to be represented unless/until I instructed otherwise.

Please direct all correspondence in this case to at the address indicated below:

Ronald J. Baron Hoffman & Baron, L.L.P. 6900 Jericho Turnpike Syosset, New York 11791

Family Name	First Given Name	Second Given Name	
Wolties-			
	Residence and Citizenship		
City of Residence	State or Country of Residence	Country of Citizenship	
Veendam	The Netherlands	The Netherlands	
	Post Office Address		
Street Address	City /	State & Zip Code or Country	
Groningenlaan 42	Verendem	9642 EJ	
Signature of Inventor	MUS	Date 27-7-2001	

NLX

Full Name of Second Inventor, if any				
Family Name	First Given Name	Second Given Name		
Meima.	Heine	Roelf		
	Residence and Citizenship			
City of Residence	State or Country of Residence	Country of Citizenship		
Borgercompanie	The Netherlands	The Netherlands		
	Post Office Address			
Street Address	City	State & Zip Code or Country		
Borgercompanie 239	Boggercompanie	9631 TJ		
Signature of Inventor	9	Date 27-7-2001		

3-00

	Full Name of Third Inventor, if a		
Family Name	First Given Name	Second Given Name	
Buwalda Pieter_		Lykle	
	Residence and Citizenship		
City of Residence	State or Country of Residence	Country of Citizenship	
Groningen The Netherlands		The Netherlands	
	Post Office Address		
Street Address	City	State & Zip Code or Country	
Mondriaanstraat 32	Groningen	9718 MJ	
Signature of Inventor	01 1/1/2	Date 27-7-2601	

NLX

NLX

# Declaration and Power of Attorney Patent Application (Design or Utility)

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name,

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled: 'Composition based on cross-linked starch and depolymerized starch suitable as gelatine replacement'

the	specification	of	which

П	is attached hereto		
x	was filed on July 9,	2001 as application serial no.	09/889.116
	and or PCT Internation	al Application number PCT/NL00/00059 and	d was amended
	on	(if applicable).	

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose to the U.S. Patent and Trademark Office all information know to me to be material to patentability as defined in 37 C.F.R.§1.56.

I hereby claim foreign priority benefits under 35 U.S.C.§119(a)-(d) or 35 U.S.C.§365(b) of any foreign application(s) for patent or inventor's certificate, or 35 U.S.C.§365(a) of any PCT International application which designated at least one country other than the United States, listed below and have also identified below any foreign application for patent or inventor's certificate of PCT International application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application(s)					
Number 99200262.6	Country EP	Day/Month/Year Filed 29-01-1999			
Number	Country	Day/Month/Year Filed			
Number	Country	Day/Month/Year Filed			

I hereby claim the benefit under 35 U.S.C. §119(e) of any United States provisional application(s) listed below:

Prior Provisional Application(s)					
Serial Number	Day/Month/Year Filing Date				
Serial Number	Day/Month/Year Filing Date				
Serial Number	Day/Month/Year Filing Date				

I hereby claim the benefit under 35 U.S.C. §120 of any United States application(s), or under 35 U.S.C. §365(c) of any PCT International application designating the United States, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of 35 U.S.C. §112, I acknowledge the duty to disclose to the U.S. Patent and Trademark Office all information known to me to be material to patentability as defined in 37 C.F.R.§1.56 which became available between the filing date of the prior application and the national or PCT International filing date of this application:

Prior U.S. or International Application(s)					
Serial Number	Day/Month/Year Filed	Status (patented, pending, abandoned)			
Serial Number	Day/Month/Year Filed	Status (patented, pending, abandoned)			
Serial Number	Day/Month/Year Filed	Status (patented, pending, abandoned)			

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements are made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under 18 U.S.C.§1001 and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.